

Figure 60A provides a schematic showing an arrangement of a target-specific invader oligonucleotide (SEQ ID NO:61) and a target-specific probe oligonucleotide (SEQ ID NO:62) bearing a 5' Cy3 label along a target nucleic acid (SEQ ID NO:63).

Figure 60B is the image generated by a fluorescence imager showing the detection of specific cleavage products generated in an invasive cleavage assay using charge reversal (*i.e.*, charge based separation of cleavage products).

Figure 61 is the image generated by a fluorescence imager which depicts the sensitivity of detection of specific cleavage products generated in an invasive cleavage assay using charge reversal.

Figure 62 depicts a first embodiment of a device for the charge-based separation of oligonucleotides.

Figure 63 depicts a second embodiment of a device for the charge-based separation of oligonucleotides.

Figure 64 shows an autoradiogram of a gel showing the results of cleavage reactions run in the presence or absence of a primer oligonucleotide; a sequencing ladder is shown as a size marker.

Figures 65a-d depict four pairs of oligonucleotides; in each pair shown, the upper arrangement of a probe annealed to a target nucleic acid lacks an upstream oligonucleotide and the lower arrangement contains an upstream oligonucleotide.

Figure 66 shows the chemical structure of several positively charged heterodimeric DNA-binding dyes.

Figure 67 is a schematic showing alternative methods for the tailing and detection of specific cleavage products in the context of the Invader™-directed cleavage assay.

Figure 68 provides a schematic drawing of a target nucleic acid with an Invader™ oligonucleotide, a miniprobe, and a stacker oligonucleotide annealed to the target.

Figure 69 provides a space-filling model of the 3-dimensional structure of the T5 5'-exonuclease.

Figure 70 provides an alignment of the amino acid sequences of several FEN-1 nucleases including the *Methanococcus jannaschii* FEN-1 protein (MJAFEN1.PRO), the *Pyrococcus furiosus* FEN-1 protein (PFUFEN1.PRO), the human FEN-1 protein (HUMFEN1.PRO), the mouse FEN-1 protein (MUSFEN1.PRO), the *Saccharomyces cerevisiae* YKL510 protein (YST510.PRO), the *Saccharomyces cerevisiae* RAD2 protein (YSTRAD2.PRO), the *Shizosaccharomyces pombe* RAD13 protein (SPORAD13.PRO), the human XPG protein (HUMXPG.PRO), the mouse XPG protein (MUSXPG.PRO), the *Xenopus laevis* XPG protein (XENXPG.PRO) and the *C. elegans* RAD2 protein (CELRAD2.PRO); portions of the amino acid sequence of some of these proteins were not shown in order to maximize the alignment between proteins. The numbers to the left of each line of sequence refers to the amino acid residue number; dashes represent gaps introduced to maximize alignment.

Figure 71 provides a schematic showing the S-33 and 11-8-0 oligonucleotides in a folded configuration; the cleavage site is indicated by the arrowhead.

DEFINITIONS

As used herein, the terms "complementary" or "complementarity" are used in reference to polynucleotides (*i.e.*, a sequence of nucleotides such as an oligonucleotide or a target nucleic acid) related by the base-pairing rules. For example, for the sequence "A-G-T," is complementary to the sequence "T-C-A." Complementarity may be "partial," in which only some of the nucleic acids' bases are matched according to the base pairing rules. Or, there may be "complete" or "total" complementarity between the nucleic acids. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, as well as detection methods which depend upon binding between nucleic acids.

The term "homology" refers to a degree of identity. There may be partial homology or complete homology. A partially identical sequence is one that is less than 100% identical to another sequence.

As used herein, the term "hybridization" is used in reference to the pairing of complementary nucleic acids. Hybridization and the strength of hybridization (*i.e.*, the strength of the association between the nucleic acids) is impacted by such factors as the degree of complementarity between the nucleic acids, stringency of the conditions involved, the T_m of the formed hybrid, and the G:C ratio within the nucleic acids.

As used herein, the term " T_m " is used in reference to the "melting temperature." The melting temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. The equation for calculating the T_m of nucleic acids is well known in the art. As indicated by standard references, a simple estimate of the T_m value may be calculated by the equation: $T_m = 81.5 + 0.41(\% G + C)$, when a nucleic acid is in aqueous solution at 1 M NaCl (*see e.g.*, Anderson and Young, Quantitative Filter Hybridization, in *Nucleic Acid Hybridization* (1985). Other references include more sophisticated computations which take structural as well as sequence characteristics into account for the calculation of T_m .

As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds, under which nucleic acid hybridizations are conducted. With "high stringency" conditions, nucleic acid base pairing will occur only between nucleic acid fragments that have a high frequency of complementary base sequences. Thus, conditions of "weak" or "low" stringency are often required when it is desired that nucleic acids which are not completely complementary to one another be hybridized or annealed together.

The term "gene" refers to a DNA sequence that comprises control and coding sequences necessary for the production of a polypeptide or precursor. The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired enzymatic activity is retained.

The term "wild-type" refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designated the "normal" or "wild-type" form of the gene. In